



International
Animal Health Products
THE AUSTRALASIAN COMPANY

CTC-ECO™ 200 GRANULAR

FEED SUPPLEMENT FOR PROPHYLACTIC USE WHEN AT RISK FROM DISEASES AND FOR TREATMENT OF DISEASES CAUSED BY MICROORGANISMS SUSCEPTIBLE TO CHLORTETRACYCLINE IN CALVES, PIGS AND POULTRY.

PRESENTATION: Brown free flowing micro-granules.

ACTIVE CONSTITUENTS: Each kg contains:

200 g/kg CHLORTETRACYCLINE HYDROCHLORIDE (as calcium complex)

PROPERTIES: CTC-ECO™ BASE 200 is the feed stable, calcium complex of chlortetracycline hydrochloride, a broad-spectrum antibiotic, highly effective against a variety of gram-positive and gram-negative pathogenic bacteria such as Mycoplasmas, Ehrlichia, Rickettsias, Chlamidias and some atypical mycobacteria and amoeba that are not controlled by other antibiotics and bacteriostatic agents.

CTC-ECO™ 200 GRANULAR is specifically designed for the inclusion in animal feed. It is non-dusty micro-granules to give maximum dispersion in feed and in addition is well absorbed. It is generally very stable in the dry form, particularly when compared to other antibiotics.

DOSAGE AND ADMINISTRATION:

DO NOT USE in lactating cows where milk or milk products may be used for human consumption.

Prophylactic dose: If calves, pigs or poultry at risk of stress such as rehousing, transport, vaccination, weaning or in conditions that predispose towards disease, CTC-ECO™ 200 GRANULAR should be supplied at prophylactic dose for periods up to 28 days. Thoroughly mix 500 g CTC-ECO™ 200 GRANULAR into 1 tonne of finished feed (equivalent to 100 ppm chlortetracycline hydrochloride).

Therapeutic dose: Thoroughly mix 2 kg CTC-ECO™ 200 GRANULAR into 1 tonne of finished feed (equivalent to 400 ppm chlortetracycline hydrochloride) and dose for not more than 7 days.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

WITHHOLDING PERIOD:

Prophylactic dose: MEAT: REMOVE ALL MEDICATED FEED, poultry – 2 days; calves & pigs – 5 days before slaughter for human consumption.

Therapeutic dose: MEAT: REMOVE ALL MEDICATED FEED, calves – 10 days; poultry & pigs – 7 days before slaughter for human consumption.

EGGS – NIL

DO NOT USE in lactating cows where milk or milk products may be used for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9672 7944 before using this product.

POISONS SCHEDULE: S4 (Prescription Animal Remedy)

REGULATORY STATUS: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY

PACK SIZE: 25 kg

Australia-Wide Toll Free: 1800 801 201

Email: info@iahp.com.au

Website: www.iahp.com.au

® The Australian/World symbol is a registered trademark of International Animal Health Products P/L.

™ CTC-ECO is a trademark of International Animal Health Products P/L.

© Copyright 2002–2021. All Rights Reserved.

01/2021