



CTC-ECO™ ORAL POWDER

WATER SOLUBLE CONTROL AND TREATMENT OF DISEASES CAUSED BY CHLORTETRACYCLINE-SUSCEPTIBLE MICROORGANISMS IN POULTRY AND PIGS.

PRESENTATION: A hygroscopic, golden-yellow crystalline powder.

ACTIVE CONSTITUENTS: Each kg contains: min. 950 g CHLORTETRACYCLINE HYDROCHLORIDE

PROPERTIES: CTC-ECO™ ORAL POWDER is a concentrated chlortetracycline hydrochloride, a broad-spectrum antibiotic. Chlortetracycline is highly effective against a variety of gram positive and gram negative bacteria such as *Mycoplasma*, *E. coli*, *Coryza*, *Salmonella*, *Staphylococcus*, *Pasteurella*, *Hexamitia*; against bacterial enteritis in poultry; and against bacterial enteritis, pneumonia and secondary infection in pigs. It also useful in Rickettsial diseases, infections produced by some *Mycoplasma*, disorders caused by *Chlamydia*, some atypical mycobacteria and amoeba that are not controlled by other antibiotics and bacteriostatic agents.

DOSAGE AND ADMINISTRATION:

Medicated water should be prepared daily. All other sources of water should be removed until the medicated water has been consumed. Medicated water should be consumed within 24 hours.

POULTRY – add to drinking water.

PIGS – add to drinking water, milk or milk replacer, or use as a drench.

In poultry or pigs at risk of stress such as transport, or in conditions that predispose to towards disease, CTC-ECO™ ORAL POWDER should be supplied at a prophylactic dose for periods up to 28 days.

Prophylactic dose: Supply daily for 3 or more days as indicated. **Poultry:** 12 mg/kg body weight. **Pigs:** 6 mg/kg body weight.

Therapeutic dose: Supply for 2 to 5 days, depending on the severity of the symptoms. Continue administration for 2 to 3 days after all symptoms have subsided. **Poultry:** 60 mg/kg body weight. **Pigs:** 25 mg/kg body weight.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may result in the need to extend the approved withholding period.

WITHHOLDING PERIOD:

Therapeutic dose: MEAT – DO NOT USE less than: Poultry and pigs – 7 days before slaughter for human consumption.

Prophylactic dose: MEAT – DO NOT USE less than: Poultry – 4 days; pigs – 5 days before slaughter for human consumption. EGGS – NIL

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9672 7944 before using this product.

POISONS SCHEDULE: S4 (Prescription Animal Remedy)

REGULATORY STATUS: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY

PACK SIZE: 200 g, 500 g, 25 kg

Australia-Wide Toll Free: 1800 801 201

Email: info@iahp.com.au

Website: www.iahp.com.au

® The Australian/World symbol is a registered trademark of International Animal Health Products P/L.

™ CTC-ECO is a trademark of International Animal Health Products P/L.

© Copyright 2001–2021. All Rights Reserved.

01/2021