



DOTECO 1000™

*AN AID IN THE PREVENTION OF COCCIDIOSIS OF BROILERS, REPLACEMENT PULLETS
AND TURKEYS.*

PRESENTATION: Cream to light brown free-flowing powder free of particulate matter.

ACTIVE CONSTITUENTS: DINITOLMIDE (3,5-DINITRO-ORTHO-TOLUAMIDE) 1000 g/kg.

PROPERTIES: DOTEKO™ 1000 is effective as an aid in the prevention of coccidiosis of broilers, replacement pullets and turkeys.

DOSE AND ADMINISTRATION:

Use continuously for up to 14 days or as directed by a Veterinarian. Consult a Veterinarian for advice relating to coccidiosis control programs.

DOTECO 1000™ should be premixed with 5-10 kg of feed and then thoroughly mixed with the bulk feed.

Broilers and Replacement Pullets: Add DOTEKO 1000™ at the rate of 125 grams/tonne of finished feed.

Turkey Growers: Add DOTEKO 1000™ at the rate of 190 grams/tonne of finished feed.

WITHHOLDING PERIODS:

MEAT: NIL.

EGGS: DO NOT USE in laying hens (chickens/turkeys) or in replacement pullets over 14 weeks of age.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established.

For advice on the ESI, contact the manufacturer on (02) 9672 7944 before using this product.

POISONS SCHEDULE: NOT APPLICABLE

REGULATORY STATUS: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY

PACK SIZES: 25 kg