



International
Animal Health Products
THE AUSTRALASIAN COMPANY

SALECO® 120 GRANULAR

GRANULAR ANTICOCIDIAL FEED ADDITIVE MICRO-GRANULE FOR BROILERS, STIMULATES GROWTH RATE AND INCREASED FEED CONVERSION OF PIGS AND BEEF CATTLE.

PRESENTATION: A beige, free flowing granules.

ACTIVE CONSTITUENTS: Each kg contains: 120 g SALINOMYCIN SODIUM

PROPERTIES: SALECO® 120 GRANULAR contains salinomycin sodium, which aids in prevention of coccidiosis in broiler chickens and in replacement birds intended for use as caged layers caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella*, their sporozoites, trophozoites and first generation schizonts. SALECO® 120 GRANULAR may be used in replacement pullets for periods up to 7 days before producing eggs and eggs products for human consumption, or as directed by a Veterinarian.

Coccidia are widely distributed in intensive livestock production. With the development of intensive methods of husbandry, coccidiosis control has become more difficult. The high contamination rates and risk of infection have led to the wide use of prophylactic medication that can be maintained through SALECO® 120 GRANULAR, especially in broiler production. Additionally, it also acts against gram-positive bacteria, mycobacteria and some filamentous fungi. SALECO® 120 GRANULAR improves the productivity by stimulating the growth rate and increasing the feed conversion efficiency of grower/finisher pigs and feedlot beef cattle.

DOSAGE AND ADMINISTRATION: FOR ORAL ADMINISTRATION IN FEED. MIX THOROUGHLY THE APPROPRIATE AMOUNT OF SALECO® 120 GRANULAR WITH FEED.

POULTRY: Mix 500g SALECO® 120 GRANULAR in 1 tonne of finished chicken feed to provide a salinomycin concentration of 60 g/tonne (60ppm). This medicated feed should be fed continuously as the sole ration.

PIGS: Mix 210 g SALECO® 120 GRANULAR in 1 tonne of complete pig grower/finisher feed to provide salinomycin concentration of 25 g/tonne (25ppm).

BEEF CATTLE: Mix 125 g SALECO® 120 GRANULAR in 1 tonne of cattle feed to provide a salinomycin concentration of 15 g/tonne (15ppm).

WITHHOLDING PERIOD: (CATTLE:) MEAT: NIL; MILK: DO NOT USE in animal producing milk for human consumption. (POULTRY:) MEAT: NIL; EGGS: DO NOT USE in birds laying eggs for human consumption or where egg products are to be used for human consumption or processing.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9672 7944 before using this product.

PRECAUTION: DO NOT exceed recommended dose rates. DO NOT feed to horses or any other animals or bird not included on the label. DO NOT use in animals producing milk for human consumption. DO NOT administer to animals receiving Tiamulin (Dynamutilin). Care should be taken when feeding concurrently with other antimicrobial products.

POISONS SCHEDULE: S6.

REGULATORY STATUS: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY.

PACK SIZE: 25 kg.

Australia-Wide Toll Free: 1800 801 201

Email: info@iahp.com.au

Website: www.iahp.com.au

© SALECO and the Australian/World symbol are registered trademarks of International Animal Health Products P/L.
© Copyright 2001-2020. All Rights Reserved.

04/2020