



International
Animal Health Products
THE AUSTRALASIAN COMPANY

WORMA DRENCH[®]

BROAD SPECTRUM ORAL ANTHELMINTIC FOR HORSES AND CATTLE (DAIRY & BEEF.)

MAY ALSO BE ADDED TO FEED

PRESENTATION: White emulsion, relatively odourless and tasteless.

ACTIVE CONSTITUENTS: OXFENDAZOLE 100 g/L

PROPERTIES: FARNAM[®] WORMA DRENCH[®] is effective against the following internal parasites:

Horses: Mature & immature stages of Bloodworm (*Strongylus vulgaris*); Large Strongyles or Redworms, including the migratory tissue stages of *S. edentatus*; Large Roundworm, Pinworms, Small Redworms (Small strongyles or Cyathostome species susceptible to oxfendazole), mature Stomach Worms (*Habronema microstoma*), as well as the harmful migrating stages of the Bloodworm (*S. vulgaris*) (55-88% activity) with corresponding reduction in arterial lesions.

Cattle: Dairy & Beef - Small Brown Stomach Worm (*Ostertagia ostertagi*) including inhibited fourth larval or hypobiotic stages, Barber's Pole Worm, Small Intestinal Worm, Stomach hairworm, Thin Necked Intestinal Worm, Hookworm, Nodule Worm, Lungworm, and Cattle Tapeworm.

WORMA DRENCH in liquid form provides an economic and convenient method of controlling major internal Roundworms in horses on studs or in large groups, as well as Roundworms and Tapeworms in cattle on studs, dairy herds, feedlots and grazing stock.

DOSE AND ADMINISTRATION:

Administer orally with accurate drenching equipment, disposable syringe, stomach tube, or by mixing thoroughly into the feed. Base the dose on the heaviest animal in the herd. Calculate body weight using scales or a weigh-band.

HORSES: Use 10 mL of WORMA DRENCH
per 100 kg body weight or as per table

0 – 50 kg: 5 mL	251 – 300 kg: 30 mL
51 – 100 kg: 10 mL	301 – 350 kg: 35 mL
101 – 150 kg: 15 mL	351 – 400 kg: 40 mL
151 – 200 kg: 20 mL	401 – 450 kg: 45 mL
201 – 250 kg: 25 mL	451 – 500 kg: 50 mL
500 kg & over: 10 mL/100 kg bodyweight.	

CATTLE: Use 5 mL of WORMA DRENCH
per 100 kg body weight or as per table

0 – 50 kg: 3 mL	351 – 400 kg: 20 mL
51 – 100 kg: 5 mL	401 – 450 kg: 23 mL
101 – 150 kg: 8 mL	451 – 500 kg: 25 mL
151 – 200 kg: 10 mL	501 – 550 kg: 28 mL
201 – 250 kg: 13 mL	551 – 600 kg: 30 mL
251 – 300 kg: 15 mL	601 – 650 kg: 33 mL
301 – 350 kg: 18 mL	

650 kg & over: 5 mL/100 kg bodyweight.

NOTE: Safe to use in pregnant, sick or debilitated animals, but always take care.

		<u>Adult horses doses</u>	<u>Foal doses</u>	<u>Adult cattle doses</u>	<u>Calf doses</u>
250 mL	pack contains	5	25	10	50
1 L	pack contains	20	100	40	200
5 L	pack contains	100	500	200	1000

WITHHOLDING PERIODS: (HORSES): MEAT: 28 days. (CATTLE): MILK: NIL; MEAT: DO NOT ADMINISTER LESS THAN 8 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 14 days before slaughter for export. The ESI on this datasheet was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on (02) 9672 7944 or the APVMA website (www.apvma.gov.au/residues/ESI.shtml).

POISONS SCHEDULE: S5

REGULATORY STATUS: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY

PACK SIZES: 250 mL, 1 L, 5 L, 25 L