



FARNAM® WORMA® PASTE FOR HORSES

BROAD SPECTRUM ORAL PASTE FOR HORSES OF ALL AGES.

PRESENTATION: Oral paste in dial-a-dose syringe. **FARNAM® WORMA® PASTE FOR HORSES** is safe for use in horses of all ages.

ACTIVE CONSTITUENTS: Each g contains:

OXFENDAZOLE	107.2 mg
PIPERAZINE (as DIHYDROCHLORIDE)	214.4 mg

PROPERTIES:

A broad spectrum oral anthelmintic administered by syringe. Controls sensitive strains of adult and immature forms of large roundworms (*Parascaris equorum*), pinworm (*Oxyuris equi*), large strongyles (*Strongylus* spp), adult stomach hairworm (*Trichonstrongylus axei*) and adult stomach worm (*Habronema microstoma*) and small strongyles (*Cyathostomes*). Also kills tissue stages of *Strongylus edentatus* and aids in the control of migrating stages of *Strongylus vulgaris* with a corresponding reduction in arterial lesions. Two doses given two days apart gives enhanced efficacy against migrating strongyles and will decrease the incidence of associated colic.

DOSAGE & ADMINISTRATION:

Dose according to bodyweight by setting the plunger ring to the required bodyweight setting. The syringe is calibrated in 50 kg divisions. A 56 g syringe contains sufficient **WORMA® PASTE** to treat a 600 kg horse. Ensure syringe is placed on the back of the horse's tongue, not between the teeth and cheeks. Depress the plunger and keep horse's head elevated and its mouth closed until the paste is swallowed. Internal parasites can develop resistance to anthelmintics. Pasture management, rotation between different types of wormer and other measures should be used to minimise the incidence of anthelmintic resistance.

Do not treat sick, debilitated or pregnant animals without seeking veterinary advice.

Note:

- Treat all horses on the property every 6-8 weeks. Treat visiting and new horses before introduction to the pasture.
- Foals may be treated every 4 weeks to control Large Roundworm.
- When rotating to other wormers, we recommend using **M^{EC}WORMA® & BOT** or **M^{EC}WORMA® & TAPE**.

MEAT WITHHOLDING PERIOD (HORSES): DO NOT ADMINISTER LATER THAN 28 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

POISONS SCHEDULE: S5 (CAUTION)

REGULATORY STATUS: **Australia:** AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY

PACK SIZE: 56 g (15 per carton and 30 per pail)

Australia-Wide Toll Free: 1800 801 201

Email: info@iahp.com.au

Website: www.iahp.com.au

® FARNAM, WORMA, M^{EC}WORMA and the Australia/World symbol are registered trademarks of International Animal Health Products P/L.

© Copyright 1997-2025. All Rights Reserved.

03/2025